

**In the Claims:**

1. (Withdrawn) A method of controlling Dipteran larvae comprising the step of introducing a larvicidally-effective amount of a combination of a strain of *Bacillus thuringiensis* subspecies *israelensis* and a strain of *Bacillus sphaericus* into an environment containing Dipteran larvae.

2. (Withdrawn) The method of claim 1 wherein said strain of *Bacillus thuringiensis* subspecies *israelensis* is non-genetically modified.

3. (Withdrawn) The method of claim 1 wherein said strain of *Bacillus sphaericus* is non-genetically modified.

4. (Withdrawn) The method of claim 1 wherein said strain of *Bacillus thuringiensis* subspecies *israelensis* is non-genetically modified and said strain of *Bacillus sphaericus* is non-genetically modified.

5. (Withdrawn) The method of claim 1 wherein said combination has from about 1:10 to about 10:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

6. (Withdrawn) The method of claim 1 wherein said combination has from about 1:3 to about 3:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

7. (Withdrawn) The method of claim 1 wherein said combination has from about 1:2 to about 2:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

8. (Withdrawn) The method of claim 1 wherein said combination has a 1:1 ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

9. (Withdrawn) The method of claim 1 wherein said Dipteran is a mosquito.

10. (Withdrawn) The method of claim 9 wherein said mosquito is selected from the group consisting of *Culex pipiens*, *Culex quinquefasciatus*, *Aedes aegypti*, *Culex tarsalis*, *Culiseta incidens*, *Anopheles freeborni* and combinations thereof.

11. (Withdrawn) A method for inhibiting larvicidal resistance in Diptera comprising the step of introducing a larvicidally-effective amount of a combination of a strain of *Bacillus thuringiensis* subspecies *israelensis* and a strain of *Bacillus sphaericus* into an environment containing Dipteran larvae.

12. (Withdrawn) The method of claim 11 wherein said strain of *Bacillus thuringiensis* subspecies *israelensis* is non-genetically modified.

13. (Withdrawn) The method of claim 11 wherein said strain of *Bacillus sphaericus* is non-genetically modified.

14. (Withdrawn) The method of claim 11 wherein said strain of *Bacillus thuringiensis* subspecies *israelensis* is non-genetically modified and said strain of *Bacillus sphaericus* is non-genetically modified.

15. (Withdrawn) The method of claim 11 wherein said combination has from about 1:10 to about 10:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

16. (Withdrawn) The method of claim 11 wherein said combination has from about 1:3 to about 3:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

17. (Withdrawn) The method of claim 11 wherein said combination has from about 1:2 to about 2:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

18. (Withdrawn) The method of claim 11 wherein said combination has a 1:1 ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

19. (Withdrawn) The method of claim 11 wherein said Diptera is *Culex*.

20. (Withdrawn) The method of claim 11 wherein larvicidal resistance is developed against *Bacillus sphaericus*.

21. (Withdrawn) The method of claim 11 wherein said Diptera is a mosquito.

22. (Withdrawn) The method of claim 21 wherein said mosquito is selected from the group consisting of *Culex pipiens*, *Culex quinquefasciatus*, *Aedes aegypti*, *Culex tarsalis*, *Culiseta incidens*, *Anopheles freeborni* and combinations thereof.

23. (Previously Presented) A composition comprising:  
a combination of an isolated, biologically pure strain of *Bacillus thuringiensis* subspecies *israelensis* and an isolated, biologically pure strain of *Bacillus sphaericus*.

24. (Previously Presented) The composition of claim 23 wherein said strain of *Bacillus thuringiensis* subspecies *israelensis* is wild-type.

25. (Previously Presented) The composition of claim 23 wherein said strain of *Bacillus sphaericus* is wild-type.

26. (Previously Presented) The composition of claim 23 wherein said strain of *Bacillus thuringiensis* subspecies *israelensis* is wild-type and said strain of *Bacillus sphaericus* is wild-type.

27. (Previously Presented) The composition of claim 23 wherein said combination has from about 1:10 to about 10:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

28. (Previously Presented) The composition of claim 23 wherein said combination has from about 1:3 to about 3:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

29. (Previously Presented) The composition of claim 23 wherein said combination has from about 1:2 to about 2:1 weight ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

30. (Previously Presented) The composition of claim 23 wherein said combination has a 1:1 ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

31. (Previously Presented) The composition of claim 23 further comprising an additional component selected from the group consisting of a surface active agent, an inert carrier, a preservative, a humectant, a feeding stimulant, an attractant, an encapsulating agent, a binder, an emulsifier, a dye, a U.V. protectant, a buffer, a drift control agent, a spray deposition aid, a free-flow agent and combinations thereof.

32. (Previously Presented) The composition of claim 23 wherein slurries of both strains are spray dried.

33. (Previously Presented) The composition of claim 32 wherein the slurries are each mixed with useful compositions and then each mixture is spray dried separately.

34. (Previously Presented) The composition of claim 32 wherein the slurries are mixed together, then preserved and then the mixture is spray dried.

35. (Previously Presented) The composition of claim 23 wherein said combination has a 4:1 ratio of *Bacillus thuringiensis* subspecies *israelensis* to *Bacillus sphaericus*.

36. (Withdrawn) A method of using the composition of claim 34 for management of resistance wherein the bioactivity of said composition are at a predetermined concentration and based on solids or biopotency, said concentration is selected from the group consisting of: 1:1, 2:1, 3:1, 1:2, 1:3, and 1:4 ratio of *Bacillus thuringiensis* subsp. *Israelensis* and *Bacillus sphaericus*, said method comprises the steps of preserving, adding stabilizers, adding surfactants, adding UV protectant, and delivering to aquatic mosquito habitats to eliminate or substantially minimize potential for build up of resistance.